

A.N. Sazonov

Optical Multicolor Observations of the SS 433=V 1343 Aql Microquasar

2009 A.N. Sazonov

Sternberg Astronomical Institute, Universitetskii pr. 13, Moscow, 119992 Russia

We report BVR photometry of the V1343 Aql= SS 433 microquasar at different phases of the 13-day orbital cycle for the 1986-1990 observing seasons. The data include five complete cycles of the 163^d precession period of the system. We obtain mean light curves and color-color diagrams with the orbital period for all intervals of precession phases. The optical component of the close binary system (CBS) fills its critical Roche lobe and loses mass on the thermal relaxation time scale. Gaseous flow show up actively in the system and activity manifestations differ substantially at different precession phases.

The collimated relativistic jets perpendicular to the plane of the disk appear to be associated with supercritical accretion onto the compact relativistic object in the massive CBS, which shows up in the shapes of the light curves at different orbital and precession phases. An analysis of color indices confirmed the earlier discovered peculiarities of the system ([1]): (1) A "disk corona" around the compact object. (2) Phase shifts between orbital light curves and different heights of light maxima for different passbands and at all phases of the 163^d precession period.

The BVR light curves of the CBS V1343 Aql= SS 433 are qualitatively similar, the depth of the primary minimum (the eclipse of the accretion disk of the system by its "normal" - star component) during the observing seasons considered was of about $0^m.65 \div 0^m.75$, $0^m.50 \div 0^m.60$, and $0^m.35 \div 0^m.45$ in the B, V, and R passbands, respectively. The light curves vary with the precession phase in what must be a manifestation of the activity of gaseous flows of the CBS and their active interaction with the "floating" accretion disk whose position in space is related with the orientation of relativistic jets.

The phase shifts of orbital light curves and different heights of light maxima confirm the asymmetry of the brightness distribution in the accretion disk and the fact that the downstream side of the AD (with respect to orbital motion) is brighter than the upstream side.

Of considerable interest and value are multicolor photometric observations of SS 433 [1], [11]), because most of the extensive V-band, and much scarcer B and R-band observations

of this unique object were made in only, mostly the V-band, filter [3]; [4]; [5]; [6];

The color variations of the star with the orbital and precession periods at Min I and Min II have been confirmed conclusively during the observing seasons considered .

[1] obtained a number of original results obtained for the first time with the AZT-11 telescope equipped with a five-color UBVRI photometer of the Helsinki University [7]; [1]; [8] with a large aperture.

The device can be used to perform synchronous polarimetric and photometric observations in five bands close to the standard bands of the Johnson's UBVRI system, which is very important for a quantitative and qualitative comparison with other observations made in the same photometric systems.

During the 1986-1990 observing period a series of five to seven outbursts occurred in each observing season (especially in 1988) with the V-band amplitudes ranging from $\sim 0^m.30$ to $\sim 0^m.50$. Optical outbursts of SS 433 had almost uniform distribution of precession phases during these observing seasons.

Observations were performed on time scales of fast variability with exposures ranging from 90^s to 180^s on time intervals from 1 to 2.5 hours during the night.

The small amplitude of B-V color index variations shows up conspicuously in the color-index diagrams. It is evidently due to the relatively small B-V color differences between the thermal continuum emitted by the bright optical star and different parts of the accretion disk of the system ([1]), because above 20000 K (B-V) colors are practically independent of temperature.

Note that the behavior of periodic light variations of SS 433 during these observing seasons qualitatively agrees with the data by other authors for the period from 1978 through 1990, suggesting a certain stability of regular light variations of the system with the periods of $163^d.34$ (the precession period, the amplitude $\Delta V \cong 0^m.75 \div 0^m.85$) and $13^d.086$ (the orbital period, the corresponding amplitude is $\Delta V \cong 0^m.75 \div 0^m.85$).

We also studied the nutation light curve of the object based on observations made in 1986-1990, i.e., the light curve of variations that remain after the subtraction of precession and orbital variations and folded with the nutation period of 6.28 d. The computed ephemeris for the maximum light of nutation variations is ([9]):

$$Max = JD2450000^d.94 + 6^d.2877E$$

[10] pointed out that one of the most salient features of the behavior of relativistic jets as a function of the phase of the nutation cycle is the sharp increase of the equivalent width W_λ of relativistic lines at the bending points of the radial-velocity curves and that this effect is especially conspicuous at phases $\sim \psi=0.00$, where jets are most inclined to the line of sight of a terrestrial observer. Moreover, at these phases variation of the angle between the line of sight and direction of the jet is little affected by the precession motion of relativistic jets and the angle considered varies mostly due to nutational oscillation of the jets. Unpredictable enhancements of line components are sometimes observed at other nutational phases (e.g., the H_α^- line). There is a viewpoint that these features are due to the nonstationary nature of the activity of SS 433 and strong variations of the mass-loss rates in relativistic jets. However, on the other hand, there may be other singular points besides the bending points on the radial-velocity curves, e.g., *self-crossing points of the jets of the gaseous stream*.

[11] also pointed out the above property of the trajectories of the particles of the gaseous stream in a CBS — the intersection (even in the celestial-mechanics approximation). The fact that ballistic trajectories of stream particles intersect may prove to be one of the factors conducive to the formation of irregularities in the gaseous jet, which result in scintillations of the brightness of the “hot spot” that forms where the jet encounters the disk-like envelope of the relativistic object of the CBS, and also in the case of self-crossing of the jet trajectory including the portions of the collimated-jet trajectory near T3 phase (more precisely, at phases $\psi = 0.03 - 0.15$).

[12]; [13] and [14] in his review point out various gaseous-flow instabilities near the compact object in the CBS. The above authors also point out that observed manifestations of gas flows include quasi-periodic oscillations of accretion rate $\dot{M}$ and the rate of change of the angular momentum, J , of the matter located in the vicinity of the secondary component of the CBS. The computed density, velocity, and temperature fields were used to determine the emission profiles of the H_β and establish that line wings form in the region near the accretor and that their broadening is determined by high velocities of the gas flow in the accreting matter and in the disk. *Gas velocities near the accretor are more than 10 times higher than the average gas velocity in the system* [13].

It was also established earlier that gas in the jet concentrates in individual clouds [16]; [15]; [17]) a pattern, which is consistent with the scenario proposed to explain

the observed data points with somewhat higher-than-average amplitude at precession phases $\psi=0.02-0.16$ (and at different orbital phases ϕ) in SS 433.

-
- [1] Shakhovskoi N.M., Sazonov A.N. // 1996, Pis'ma Astron. Zh"/", т.22, с. 8, с.580-586.
 - [11] Sazonov (A.N. Sazonov, Трехцветная фотометрия SS 433 = V1343 Aql в 1986-87 гг.) // 1988, Astron. Tsirk., no. 1531.
 - [3] (Cherepashchuk A.M. // 1981a, SS 433 as an eclipsing binary). Month. Not. Ray. Astron. Soc. **194**, 761 .
 - [4] (Kemp J.C., Henson G.D., Krauss D.I. et al.) // 1986, Astrophys.J., v.305, p.805.
 - [5] Cherepashchuk A.M., Yarikov S.F. // 1991, Письма в "Астрон. журн.", V.17, P.605.
 - [6] Irmambetova T.P. // 1997, Pis'ma "Astron. Zhurn.", V.23, no. 5, P.341-349.
 - [7] (Piirola V.) // 1973, Astron. and Astrophys. v.27. p.383.
 - [8] (Korhonen T., Piirola V., Reiz A.) // 1983, ESO Messenger. November. P.38.
 - [9] Goranskii V.P., Esipov V.F., Cherepashchuk A.M. // 1998, Astron. Zhurn., V.75, no. 2, P. 240.
 - [10] Kopylov I.M., Kumaigorodskaya R.N., Somov N.N., Somova T.A., Fabrika S.N. // 1987, Astron. Zhurn., V.64, vyp.4, P.786.
 - [11] (A.N. Sazonov, N.I. Shakura // Comm. Special Astrophys. Obs. of the Acad. Sci. of the USSR, Vyp. 64, September 19- 24, 1988).
 - [12] Bisikalo D.V., Boyarchuk A.A., Kuznetsov O.A., Popov Yu.P., Chechetkin V.M. // 1994, Astron. Zhurn., V.71, P.560.
 - [13] Bisikalo D.V., Boyarchuk A.A., Kuznetsov O.A., Chechetkin V.M. // 1996, Astron Zhurn., V.73, P.717.
 - [14] Lubow S.H. // 1993, The Realm of Interacting Binary Stars/Eds Sahade J., McCluskey J.Jr., Kondo Y. Dordrecht:Kluwer, p.25.
 - [15] Murdin P., Clark D.H., Martin P.G. // 1980, Monthly Notices Roy. Astron. Soc. v.193, p.135.
 - [16] Shklovskii I.S. // 1981, Astron. Zhurn., V.58, P. 554.
 - [17] Grandi S.A., Stone R.P.S. // 1982, Publs. Astron. Soc. Pacific., v.94, p.80.